

ALI MUHAMMAD

**On inclusion relationships  
of certain subclasses of meromorphic functions  
involving integral operator**

ABSTRACT. In this paper, we introduce some subclasses of meromorphic functions in the punctured unit disc. Several inclusion relationships and some other interesting properties of these classes are discussed.

**1. Introduction.** Let  $\mathcal{M}$  denote the class of functions  $f(z)$  of the form

$$(1.1) \quad f(z) = \frac{1}{z} + \sum_{k=0}^{\infty} a_k z^k,$$

which are analytic in the punctured open unit disc

$$E^* = \{z : z \in \mathbb{C} \text{ and } 0 < |z| < 1\} = E \setminus \{0\}.$$

If  $f(z)$  is given by (1.1) and  $g(z)$  is given by

$$(1.2) \quad g(z) = \frac{1}{z} + \sum_{k=0}^{\infty} b_k z^k,$$

---

2000 *Mathematics Subject Classification.* 30C45, 30C50.

*Key words and phrases.* Meromorphic functions, functions with bounded boundary and bounded radius rotation, quasi-convex functions, close-to-convex functions, generalized hypergeometric functions, functions with positive real part, Hadamard product (or convolution), linear operators.

we define the Hadamard product (or convolution) of  $f(z)$  and  $g(z)$  by

$$(1.3) \quad (f \star g)(z) = \frac{1}{z} + \sum_{k=0}^{\infty} a_k b_k z^k = (g \star f)(z) \quad (z \in E).$$

Let  $P_k(\rho)$  be the class of functions  $p(z)$  analytic in  $E$  with  $p(0) = 1$  and

$$(1.4) \quad \int_0^{2\pi} \left| \frac{\operatorname{Re} p(z) - \rho}{1 - \rho} \right| d\theta \leq k\pi, \quad z = re^{i\theta},$$

where  $k \geq 2$  and  $0 \leq \rho < 1$ . This class was introduced by Padmanbhan et al. in [16]. We note that  $P_k(0) = P_k$ , see [17],  $P_2(\rho) = P(\rho)$ , the class of analytic functions with positive real part greater than  $\rho$  and  $P_2(0) = P$ , the class of functions with positive real part. From (1.4) we can easily deduce that  $p(z) \in P_k(\rho)$  if and only if, there exists  $p_1(z), p_2(z) \in P(\rho)$  such that for  $z \in E$ ,

$$(1.5) \quad p(z) = \left( \frac{k}{4} + \frac{1}{2} \right) p_1(z) - \left( \frac{k}{4} - \frac{1}{2} \right) p_2(z).$$

In recent years, several families of integral operators and differential operators were introduced using Hadamard product (or convolution). For example, we choose to mention the Rushcheweyh derivative [18], the Carlson–Shaffer operator [1], the Dziok–Srivastava operator [4], the Noor integral operator [14], also see [3, 5, 6, 11]. Motivated by the work of N. E. Cho and K. I. Noor [2, 9], we introduce a family of integral operators defined on the space of meromorphic functions in the class  $\mathcal{M}$ . By using these integral operators, we define several subclasses of meromorphic functions and investigate various inclusion relationships and some other properties for the meromorphic function classes introduced here.

For complex parameters  $\alpha_1, \dots, \alpha_q$  and  $\beta_1, \dots, \beta_s$  ( $\beta_j \in \mathbb{C} \setminus \mathbb{Z}_0^-, j = 1, \dots, s$ ;  $\mathbb{Z}_0^- = \{0, -1, -2, \dots\}$ ) we now define the function  $\phi(\alpha_1, \dots, \alpha_q; \beta_1, \dots, \beta_s; z)$  by

$$\phi(\alpha_1, \dots, \alpha_q; \beta_1, \dots, \beta_s; z) = \frac{1}{z} + \sum_{k=0}^{\infty} \frac{(\alpha_1)_{k+1} \dots (\alpha_q)_{k+1}}{(\beta_1)_{k+1} \dots (\beta_s)_{k+1} (k+1)!} z^k,$$

( $q \leq s + 1$ ;  $s \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$ ;  $\mathbb{N} = \{1, 2, \dots\}$ ;  $z \in E$ ),

where  $(v)_k$  is the Pochhammer symbol (or shifted factorial) defined in (terms of the Gamma function) by

$$(v)_k = \frac{\Gamma(v+k)}{\Gamma(v)} = \begin{cases} 1 & \text{if } k = 0 \text{ and } v \in \mathbb{C} \setminus \{0\} \\ v(v+1) \dots (v+k-1) & \text{if } k \in \mathbb{N} \text{ and } v \in \mathbb{C}. \end{cases}$$

Now we introduce the following operator

$$I_{\mu}^P(\alpha_1, \dots, \alpha_q, \beta_1, \dots, \beta_s) : \mathcal{M} \longrightarrow \mathcal{M}$$

as follows:

Let  $F_{\mu,p}(z) = \frac{1}{z} + \sum_{k=0}^{\infty} \left(\frac{k+\mu+1}{\mu}\right)^p z^k$ ,  $p \in \mathbb{N}_0$ ,  $\mu \neq 0$  and let  $F_{\mu,p}^{-1}(z)$  be defined such that

$$F_{\mu,p}(z) * F_{\mu,p}^{-1}(z) = \phi(\alpha_1, \dots, \alpha_q; \beta_1, \dots, \beta_s; z).$$

Then

$$(1.6) \quad I_{\mu}^p(\alpha_1, \dots, \alpha_q, \beta_1, \dots, \beta_s)f(z) = F_{\mu,p}^{-1}(z) * f(z).$$

From (1.6) it can be easily seen

$$(1.7) \quad \begin{aligned} & I_{\mu}^p(\alpha_1, \dots, \alpha_q, \beta_1, \dots, \beta_s)f(z) \\ &= \frac{1}{z} + \sum_{k=0}^{\infty} \left(\frac{\mu}{k+\mu+1}\right)^p \frac{(\alpha_1)_{k+1} \dots (\alpha_q)_{k+1}}{(\beta_1)_{k+1} \dots (\beta_s)_{k+1} (k+1)!} a_k z^k. \end{aligned}$$

For conveniences, we shall henceforth denote

$$(1.8) \quad I_{\mu}^p(\alpha_1, \dots, \alpha_q, \beta_1, \dots, \beta_s)f(z) = I_{\mu}^p(\alpha_1, \beta_1)f(z).$$

For the choices of the parameters  $p = 0$ ,  $q = 2$ ,  $s = 1$ , the operator  $I_{\mu}^p(\alpha_1, \beta_1)f(z)$  is reduced to an operator by N. E. Cho and K. I. Noor [2] and K. I. Noor [9] and when  $p = 0$ ,  $q = 2$ ,  $s = 1$ ,  $\alpha_1 = \lambda$ ,  $\alpha_2 = 1$ ,  $\beta_1 = (n + 1)$ , the operator  $I_{\mu}^p(\alpha_1, \beta_1)f(z)$  is reduced to an operator recently introduced by S.-M. Yuan et al. in [20].

It can be easily verified from the above definition of the operator  $I_{\mu}^p(\alpha_1, \beta_1)$  that

$$(1.9) \quad z(I_{\mu}^{p+1}(\alpha_1, \beta_1)f(z))' = \mu I_{\mu}^p(\alpha_1, \beta_1)f(z) - (\mu + 1)I_{\mu}^{p+1}(\alpha_1, \beta_1)f(z)$$

and

$$(1.10) \quad z(I_{\mu}^p(\alpha_1, \beta_1)f(z))' = \alpha_1 I_{\mu}^p(\alpha_1 + 1, \beta_1)f(z) - (\alpha_1 + 1)I_{\mu}^p(\alpha_1, \beta_1)f(z).$$

By using the operator  $I_{\mu}^p(\alpha_1, \beta_1)$ , we now introduce the following subclasses of meromorphic functions:

**Definition 1.1** ([9]). A function  $f \in \mathcal{M}$  is said to belong to the class  $MR_k(\gamma)$  for  $z \in E^*$ ,  $0 \leq \gamma < 1$ ,  $k \geq 2$ , if and only if

$$-\frac{zf'(z)}{f(z)} \in P_k(\gamma),$$

and  $f \in MV_k(\gamma)$ , for  $z \in E^*$ ,  $0 \leq \gamma < 1$ ,  $k \geq 2$ , if and only if

$$-\frac{(zf(z))'}{f'(z)} \in P_k(\gamma).$$

We call  $f \in MR_k(\gamma)$  a meromorphic function with bounded radius rotation of order  $\gamma$  and  $f \in MV_k(\gamma)$  a meromorphic function with bounded boundary rotation.

**Definition 1.2.** Let  $f \in \mathcal{M}$ ,  $0 \leq \gamma < 1$ ,  $k \geq 2$ ,  $z \in E^*$ . Then

$$f \in MR_{k,\mu}^p(\alpha_1, \beta_1, \gamma) \text{ if and only if } I_\mu^p(\alpha_1, \beta_1)f \in MR_k(\gamma).$$

Also

$$f \in MV_{k,\mu}^p(\alpha_1, \beta_1, \gamma) \text{ if and only if } I_\mu^p(\alpha_1, \beta_1)f \in MV_k(\gamma), \quad z \in E^*.$$

We observe that, for  $z \in E^*$ ,

$$f \in MV_{k,\mu}^p(\alpha_1, \beta_1, \gamma) \Leftrightarrow -zf' \in MR_{k,\mu}^p(\alpha_1, \beta_1, \gamma).$$

**Definition 1.3.** Let  $\lambda \geq 0$ ,  $f \in \mathcal{M}$ ,  $p \in \mathbb{N}_0$ ,  $0 \leq \gamma, \rho < 1$ ,  $\mu > 0$  and  $z \in E^*$ . Then  $f \in B_{k,\mu}^{\lambda,p}(\alpha_1, \beta_1, \gamma, \rho)$ , if and only if there exists a function  $g \in MV_{2,\mu}^p(\alpha_1, \beta_1, \gamma)$ , such that

$$\left\{ (1 - \lambda) \frac{(I_\mu^p(\alpha_1, \beta_1)f(z))'}{(I_\mu^p(\alpha_1, \beta_1)g(z))'} + \lambda \left[ -\frac{(z(I_\mu^p(\alpha_1, \beta_1)f(z)))'}{(I_\mu^p(\alpha_1, \beta_1)g(z))'} \right] \right\} \in P_k(\rho).$$

In particular, for  $\lambda = 0 = p$ ,  $k = q = \mu = 2$  and  $s = 1$ , we obtain the class of meromorphic close-to-convex function, see [7], see also K. I. Noor [9]. For  $\lambda = 1$ ,  $p = 0$ ,  $k = q = \mu = 2$ ,  $s = 1$ , we have the class of meromorphic quasi-convex functions defined for  $z \in E^*$ . We note that the class  $C^*$  of quasi-convex univalent functions, analytic in  $E$ , was first introduced and studied in [12], see also [13, 15].

In order to establish our main results, we need the following lemma, which is properly known as the Miller–Mocanu Lemma.

**Lemma 1.1** ([8]). *Let  $u = u_1 + iu_2$ ,  $v = v_1 + iv_2$  and  $\Psi(u, v)$  be a complex valued function satisfying the conditions:*

- (i)  $\Psi(u, v)$  is continuous in a domain  $D \subset \mathbb{C}^2$ ,
- (ii)  $(1, 0) \in D$  and  $\operatorname{Re} \Psi(1, 0) > 0$ ,
- (iii)  $\operatorname{Re} \Psi(iu_2, v_1) \leq 0$ , whenever  $(iu_2, v_1) \in D$  and  $v_1 \leq -\frac{1}{2}(1 + u_2^2)$ .

*If  $h(z) = 1 + c_1z + c_2z^2 + \dots$  is a function analytic in  $E$  such that  $(h(z), zh'(z)) \in D$  and  $\operatorname{Re}(\Psi(h(z), zh'(z))) > 0$  for  $z \in E$ , then  $\operatorname{Re} h(z) > 0$  in  $E$ .*

## 2. Main results.

**Theorem 2.1.** *Let  $\operatorname{Re} \alpha_1 > 0$ ,  $\mu > 0$  and  $0 \leq \gamma < 1$ . Then*

$$MR_{k,\mu}^p(\alpha_1 + 1, \beta_1, \gamma) \subset MR_{k,\mu}^p(\alpha_1, \beta_1, \rho) \subset MR_{k,\mu}^{p+1}(\alpha_1, \beta_1, \eta).$$

**Proof.** We prove the first part of the Theorem 2.1 and the second part follows by using similar techniques. Let

$$f \in MR_{k,\mu}^p(\alpha_1 + 1, \beta_1, \gamma), \quad z \in E^*$$

and set

$$(2.1) \quad -\frac{(I_\mu^p(\alpha_1, \beta_1)f(z))'}{(I_\mu^p(\alpha_1, \beta_1)f(z))} = \left(\frac{k}{4} + \frac{1}{2}\right) h_1(z) - \left(\frac{k}{4} - \frac{1}{2}\right) h_2(z) = H(z).$$

Simple computation together with (2.1) and (1.10) yields

$$(2.2) \quad -\frac{(I_\mu^p(\alpha_1 + 1, \beta_1)f(z))'}{(I_\mu^p(\alpha_1 + 1, \beta_1)f(z))} = \left[ H(z) + \frac{zH'(z)}{-H(z) + \alpha_1 + 1} \right] \in P_k(\gamma), \quad z \in E.$$

Let

$$\Phi_{\alpha_1}(z) = \frac{1}{\alpha_1 + 1} \left[ \frac{1}{z} + \sum_{k=0}^{\infty} z^k \right] + \frac{\alpha_1}{\alpha_1 + 1} \left[ \frac{1}{z} + \sum_{k=0}^{\infty} kz^k \right],$$

then

$$(2.3) \quad \begin{aligned} H(z) * z\Phi_{\alpha_1}(z) &= \left[ H(z) + \frac{zH'(z)}{-H(z) + \alpha_1 + 1} \right] \\ &= \left( \frac{k}{4} + \frac{1}{2} \right) \left[ h_1(z) + \frac{zh_1'(z)}{-h_1(z) + \alpha_1 + 1} \right] \\ &\quad - \left( \frac{k}{4} - \frac{1}{2} \right) \left[ h_2(z) + \frac{zh_2'(z)}{-h_2(z) + \alpha_1 + 1} \right]. \end{aligned}$$

Since  $f \in MR_{k,\mu}^p(\alpha_1 + 1, \beta_1, \gamma)$ , it follows from (2.2) and (2.3) that

$$\left[ h_i(z) + \frac{zh_i'(z)}{-h_i(z) + \alpha_1 + 1} \right] \in P(\gamma), \quad i = 1, 2, \quad z \in E.$$

Let  $h_i(z) = (1 - \rho)p_i(z) + \rho$ . Then

$$\left\{ (1 - \rho)p_i(z) + \rho - \gamma + \frac{(1 - \rho)zp_i'(z)}{-(1 - \rho)p_i(z) - \rho + \alpha_1 + 1} \right\} \in P, \quad z \in E.$$

We shall show that  $p_i(z) \in P, i = 1, 2$ .

We form the functional  $\Psi(u, v)$  by taking  $u = u_1 + iu_2 = p_i(z), v = v_1 + iv_2 = zp_i'(z)$ . The first two conditions of Lemma 1.1 can be easily verified. We need to verify condition (iii) as follows:

$$\Psi(u, v) = \left\{ (1 - \rho)u + \rho - \gamma + \frac{(1 - \rho)v}{-(1 - \rho)u - \rho + \alpha_1 + 1} \right\},$$

implies that

$$\operatorname{Re} \Psi(iu_2, v_1) = \rho - \gamma + \frac{(1 - \rho)(\alpha_1 + 1 - \rho)v_1}{(1 - \rho)^2u_2^2 + (-\rho + \alpha_1 + 1)^2}.$$

By taking  $v_1 \leq -\frac{1}{2}(1 + u_2^2)$ , we have

$$\operatorname{Re} \Psi(iu_2, v_1) \leq \frac{A + Bu_2^2}{2C},$$

where

$$\begin{aligned} A &= 2(\rho - \gamma)(\alpha_1 + 1 - \rho)^2 - (1 - \rho)(\alpha_1 + 1 - \rho), \\ B &= 2(\rho - \gamma)(1 - \rho)^2 - (1 - \rho)(\alpha_1 + 1 - \rho), \\ C &= (\alpha_1 + 1 - \rho)^2 + (1 - \rho)^2u_2^2 > 0. \end{aligned}$$

We note that  $\operatorname{Re} \Psi(iu_2, v_1) \leq 0$  if and only if  $A \leq 0$  and  $B \leq 0$ . From  $A \leq 0$ , we obtain

$$(2.4) \quad \rho = \frac{1}{4} \left\{ (3 + 2\alpha_1 + 2\gamma) - \sqrt{(3 + 2\alpha_1 + 2\gamma)^2 - 8} \right\},$$

and  $B \leq 0$  gives us  $0 \leq \rho < 1$ .

Now using Lemma 1.1, we see that  $p_i(z) \in P$  for  $z \in E$ ,  $i = 1, 2$  and hence  $f \in MR_{k,\mu}^p(\alpha_1, \beta_1, \rho)$  with  $\rho$  given by (2.4).  $\square$

In particular, we note that

$$\rho = \frac{1}{4} \left\{ (3 + 2\alpha_1) - \sqrt{(12\alpha_1 + 4\alpha_1^2) + 1} \right\}.$$

**Theorem 2.2.** *Let  $\operatorname{Re} \alpha_1, \mu > 0$ . Then*

$$MV_{k,\mu}^p(\alpha_1 + 1, \beta_1, \gamma) \subset MV_{k,\mu}^p(\alpha_1, \beta_1, \rho) \subset MV_{k,\mu}^{p+1}(\alpha_1, \beta_1, \eta).$$

**Proof.** We observe that

$$\begin{aligned} f(z) \in MV_{k,\mu}^p(\alpha_1 + 1, \beta_1, \gamma) &\Leftrightarrow -zf'(z) \in MR_{k,\mu}^p(\alpha_1 + 1, \beta_1, \gamma) \\ &\Rightarrow -zf'(z) \in MR_{k,\mu}^p(\alpha_1, \beta_1, \rho) \\ &\Leftrightarrow f(z) \in MV_{k,\mu}^p(\alpha_1, \beta_1, \rho), \end{aligned}$$

where  $\rho$  is given by (2.4).

The second part can be proved by means of similar arguments.  $\square$

**Theorem 2.3.** *Let  $\operatorname{Re} \alpha_1, \mu > 0$ . Then*

$$B_{k,\mu}^{\lambda,p}(\alpha_1 + 1, \beta_1, \gamma_1, \rho_1) \subset B_{k,\mu}^{\lambda,p}(\alpha_1, \beta_1, \gamma_2, \rho_2) \subset B_{k,\mu}^{\lambda,p+1}(\alpha_1, \beta_1, \gamma_3, \rho_3),$$

where  $\gamma_i = \gamma_i(\rho_i, \mu)$ ,  $i = 1, 2, 3$  are given in the proof.

**Proof.** We prove the first inclusion of this result and the other part follows along similar lines.

Let  $f \in B_{k,\mu}^{\lambda,p}(\alpha_1 + 1, \beta_1, \gamma_1, \rho_1)$ . Then by Definition 1.3, there exists a function  $g \in MV_{2,\mu}^p(\alpha_1 + 1, \beta_1, \gamma_1)$  such that

$$(2.5) \quad \left\{ (1 - \lambda) \frac{(I_\mu^p(\alpha_1 + 1, \beta_1)f(z))'}{(I_\mu^p(\alpha_1 + 1, \beta_1)g(z))'} + \lambda \left[ -\frac{(z(I_\mu^p(\alpha_1 + 1, \beta_1)f(z)))'}{(I_\mu^p(\alpha_1 + 1, \beta_1)g(z))'} \right] \right\} \in P_k(\rho_1).$$

Set

$$(2.6) \quad h(z) = \left\{ (1 - \lambda) \frac{(I_\mu^p(\alpha_1, \beta_1)f(z))'}{(I_\mu^p(\alpha_1, \beta_1)g(z))'} + \lambda \left[ -\frac{(z(I_\mu^p(\alpha_1, \beta_1)f(z)))'}{(I_\mu^p(\alpha_1, \beta_1)g(z))'} \right] \right\},$$

where  $h(z)$  is an analytic function in  $E$  with  $h(0) = 1$ .

Now,  $g \in MV_{2,\mu}^p(\alpha_1 + 1, \beta_1, \gamma_1) \subset MV_{2,\mu}^p(\alpha_1, \beta_1, \gamma_2)$ , where  $\gamma_2$  is given by the equation

$$(2.7) \quad 2\gamma_2^2 + (3 + 2\alpha_1 - 2\gamma_1)\gamma_2 - \{2\gamma_1(1 + \alpha_1) + 1\} = 0.$$

Therefore,

$$q(z) = -\frac{(zI_\mu^p(\alpha_1, \beta_1)g(z))'}{(I_\mu^p(\alpha_1, \beta_1)g(z))'} \in P(\gamma_2), \quad z \in E.$$

By using (1.10), (2.5), (2.6) and (2.7), we have

$$(2.8) \quad \left\{ h(z) + \frac{\lambda zh'(z)}{-q(z) + \alpha_1 + 1} \right\} \in P_k(\rho_1), \quad q(z) \in P(\gamma_2), \quad z \in E.$$

With

$$h(z) = \left( \frac{k}{4} + \frac{1}{2} \right) [(1 - \rho_2)h_1(z) + \rho_2] - \left( \frac{k}{4} - \frac{1}{2} \right) [(1 - \rho_2)h_2(z) + \rho_2],$$

(2.8) can be written as

$$\begin{aligned} & \left( \frac{k}{4} + \frac{1}{2} \right) \left\{ (1 - \rho_2)h_1(z) + \rho_2 + \frac{(1 - \rho_2)\lambda zh_1'(z)}{-q(z) + \alpha_1 + 1} \right\} \\ & - \left( \frac{k}{4} - \frac{1}{2} \right) \left\{ (1 - \rho_2)h_2(z) + \rho_2 + \frac{(1 - \rho_2)\lambda zh_2'(z)}{-q(z) + \alpha_1 + 1} \right\}, \end{aligned}$$

where

$$\left\{ (1 - \rho_2)h_i(z) + \rho_2 + \frac{(1 - \rho_2)\lambda zh_i'(z)}{-q(z) + \alpha_1 + 1} \right\} \in P(\rho_1), \quad z \in E, \quad i = 1, 2.$$

That is

$$\left\{ (1 - \rho_2)h_i(z) + \rho_2 - \rho_1 + \frac{(1 - \rho_2)\lambda zh_i'(z)}{-q(z) + \alpha_1 + 1} \right\} \in P, \quad z \in E, \quad i = 1, 2.$$

We form the functional  $\Psi(u, v)$  by choosing  $u = u_1 + iu_2 = h_i(z)$ ,  $v = v_1 + iv_2 = zh_i'(z)$ , and

$$\Psi(u, v) = \left\{ (1 - \rho_2)u + \rho_2 - \rho_1 + \frac{(1 - \rho_2)\lambda v}{-q(z) + \alpha_1 + 1} \right\}, \quad (q = q_1 + iq_2).$$

The first two conditions of Lemma 1.1 are clearly satisfied. We verify (iii), with  $v_1 \leq -\frac{1}{2}(1 + u_2^2)$  as follows:

$$\begin{aligned} \operatorname{Re} \Psi(iu_2, v_1) &= \rho_2 - \rho_1 + \operatorname{Re} \left\{ \frac{\lambda(1 - \rho_2)v_1\{(-q_1 + \alpha_1 + 1) + iq_2\}}{(-q_1 + \alpha_1 + 1)^2 + q_2^2} \right\} \\ &\leq \frac{2(\rho_2 - \rho_1)|-q + \alpha_1 + 1|^2 - \lambda(1 - \rho_2)\{(-q_1 + \alpha_1 + 1)(1 + u_2^2)\}}{2|-q + \alpha_1 + 1|^2} \\ &= \frac{A + Bu_2^2}{2C} \leq 0, \end{aligned}$$

if  $A \leq 0$  and  $B \leq 0$ , where

$$\begin{aligned} A &= 2(\rho_2 - \rho_1) |-q + \alpha_1 + 1|^2 - \lambda(1 - \rho_2) \{(-q_1 + \alpha_1 + 1)\}, \\ B &= -\lambda(1 - \rho_2) \{(-q_1 + \alpha_1 + 1)\} \leq 0, \\ C &= |-q + \alpha_1 + 1|^2 > 0. \end{aligned}$$

From  $A \leq 0$ , we obtain

$$\rho_2 = \frac{2\rho_1 |-q + \alpha_1 + 1|^2 + \lambda \operatorname{Re}(-q(z) + \alpha_1 + 1)}{2 |-q + \alpha_1 + 1|^2 + \lambda \operatorname{Re}(-q(z) + \alpha_1 + 1)}.$$

Hence, using Lemma 1.1, it follows that  $h(z)$ , defined by (2.6), belongs to  $P_k(\rho_2)$  and thus  $f \in B_{k,\mu}^{\lambda,p}(\alpha_1, \beta_1, \gamma_2, \rho_2)$  for  $z \in E^*$ . This completes the proof of the first part. The second part of this result can be obtained by using similar techniques and the relation (1.9).  $\square$

**Theorem 2.4.** *Let  $\operatorname{Re} \alpha_1, \mu > 0$ . Then*

- (i)  $B_{k,\mu}^{\lambda,p}(\alpha_1, \beta_1, \gamma, \rho) \subset B_{k,\mu}^{0,p}(\alpha_1, \beta_1, \gamma, \rho_4)$ .
- (ii)  $B_{k,\mu}^{\lambda_1,p}(\alpha_1, \beta_1, \gamma, \rho) \subset B_{k,\mu}^{\lambda_2,p}(\alpha_1, \beta_1, \gamma, \rho)$ , for  $0 \leq \lambda_2 < \lambda_1$ .

**Proof.** (i). Let

$$h(z) = \frac{(I_\mu^p(\alpha_1, \beta_1)f(z))'}{(I_\mu^p(\alpha_1, \beta_1)g(z))'}$$

$h(z)$  is analytic in  $E$  and  $h(0) = 1$ . Then

$$\begin{aligned} (2.10) \quad & \left\{ (1 - \lambda) \frac{(I_\mu^p(\alpha_1, \beta_1)f(z))'}{(I_\mu^p(\alpha_1, \beta_1)g(z))'} + \lambda \left[ -\frac{(z(I_\mu^p(\alpha_1, \beta_1)f(z))')'}{(I_\mu^p(\alpha_1, \beta_1)g(z))'} \right] \right\} \\ & = h(z) + \lambda \frac{zh'(z)}{-h_0(z)}, \end{aligned}$$

where

$$h_0(z) = -\frac{(z(I_\mu^p(\alpha_1, \beta_1)f(z))')'}{(I_\mu^p(\alpha_1, \beta_1)g(z))'} \in P(\gamma).$$

Since  $f \in B_{k,\mu}^{\lambda,p}(\alpha_1, \beta_1, \gamma, \rho)$ , it follows that

$$\left[ h(z) + \lambda \frac{zh'(z)}{-h_0(z)} \right] \in P_k(\rho), \quad h_0 \in P(\gamma), \quad \text{for } z \in E.$$

Let

$$h(z) = \left( \frac{k}{4} + \frac{1}{2} \right) h_1(z) - \left( \frac{k}{4} - \frac{1}{2} \right) h_2(z).$$

Thus (2.10) implies that

$$\left[ h_i(z) + \lambda \frac{zh'_i(z)}{-h_0(z)} \right] \in P(\rho), \quad z \in E, \quad i = 1, 2.$$



and using similar techniques, together with Lemma 1.1, it follows that  $h_i(z) \in P(\rho_4)$ ,  $i = 1, 2$ , where

$$\rho_4 = \frac{2\rho |h_0(z)|^2 + \lambda \operatorname{Re} h_0(z)}{2|h_0(z)|^2 + \lambda \operatorname{Re} h_0(z)}.$$

Therefore  $h(z) \in P_k(\rho_4)$ , and  $f \in B_{k,\mu}^{0,p}(\alpha_1, \beta_1, \gamma, \rho_4)$ , for  $z \in E^*$ . In particular, it can be shown that  $h_i(z) \in P(\rho)$ ,  $i = 1, 2$ . Consequently  $h \in P_k(\rho)$  and  $f \in B_{k,\mu}^{0,p}(\alpha_1, \beta_1, \gamma, \rho)$  in  $E^*$ .

For  $\lambda_2 = 0$ , we have part (i). Therefore, we let  $\lambda_2 > 0$  and  $f \in B_{k,\mu}^{\lambda_1,p}(\alpha_1, \beta_1, \gamma, \rho)$ . There exist two functions  $H_1(z), H_2(z) \in P_k(\rho)$  such that

$$\left\{ (1 - \lambda_1) \frac{(I_\mu^p(\alpha_1 + 1, \beta_1)f(z))'}{(I_\mu^p(\alpha_1 + 1, \beta_1)g(z))'} + \lambda_1 \left[ - \frac{(z(I_\mu^p(\alpha_1 + 1, \beta_1)f(z)))'}{(I_\mu^p(\alpha_1 + 1, \beta_1)g(z))'} \right] \right\} = H_1(z)$$

$$\frac{(I_\mu^p(\alpha_1 + 1, \beta_1)f(z))'}{(I_\mu^p(\alpha_1 + 1, \beta_1)g(z))'} = H_2(z),$$

where  $g(z) \in MV_{2,\mu}^p(\alpha_1, \beta_1, \gamma)$ .

Now

$$(2.11) \quad \left\{ (1 - \lambda_2) \frac{(I_\mu^p(\alpha_1 + 1, \beta_1)f(z))'}{(I_\mu^p(\alpha_1 + 1, \beta_1)g(z))'} + \lambda_2 \left[ - \frac{(z(I_\mu^p(\alpha_1 + 1, \beta_1)f(z)))'}{(I_\mu^p(\alpha_1 + 1, \beta_1)g(z))'} \right] \right\}$$

$$= \frac{\lambda_2}{\lambda_1} H_1(z) + \left( 1 - \frac{\lambda_2}{\lambda_1} \right) H_2(z).$$

Since the class  $P_k(\rho)$  is convex, see [10], it follows that the right hand side of (2.11) belongs to  $P_k(\rho)$  and this shows that  $f \in B_{k,\mu}^{\lambda_2,p}(\alpha_1, \beta_1, \gamma, \rho)$  for  $z \in E^*$ . This completes the proof.  $\square$

**Inclusion properties involving the integral operator  $F_c$ .** Consider the operator  $F_c$ , defined by

$$(2.12) \quad F_c(f)(z) = \frac{c}{z^{c+1}} \int_0^z t^c f(t) dt \quad (f \in \mathcal{M}; c > 0).$$

From the Definition of  $F_c$  defined by (2.12), we observe that

$$(2.13) \quad z((I_\mu^p(\alpha_1, \beta_1)F_c f(z))') = c(I_\mu^p(\alpha_1, \beta_1)f(z)) - (c+1)(I_\mu^p(\alpha_1, \beta_1)F_c f(z)).$$

Using (2.12), (2.13) with similar arguments as used earlier, we can prove the following theorem.

**Theorem 2.5.** *Let  $f \in MR_{k,\mu}^p(\alpha_1, \beta_1, \gamma)$  or  $f \in MV_{k,\mu}^p(\alpha_1, \beta_1, \gamma)$  or  $f \in B_{k,\mu}^{\lambda,p}(\alpha_1, \beta_1, \gamma, \rho)$ , for  $z \in E$ . Then  $F_c(f)$  defined by (2.12) is also in the same class for  $z \in E^*$ .*

**Acknowledgement.** I am thankful for the valuable suggestions of referee which improved this paper.

#### REFERENCES

- [1] Carlson, B. C., Shaeffer, B. D., *Starlike and prestarlike hypergeometric functions*, SIAM J. Math. Anal. **15** (1984), no. 4, 737–745.
- [2] Cho, N. E., Noor, K. I., *Inclusion properties for certain classes of meromorphic functions associated with Choi–Saigo–Srivastava operator*, J. Math. Anal. Appl. **320** (2006), no. 2, 779–786.
- [3] Cho, N. E., Kwon, O. S. and Srivastava, H. M., *Inclusion relationships for certain subclasses of meromorphic functions associated with a family of multiplier transformations*, Integral Transforms Spec. Funct. **16** (2005), no. 8, 647–659.
- [4] Dziok, J., Srivastava, H. M., *Classes of analytic functions associated with the generalized hypergeometric function*, Appl. Math. Comput. **103** (1999), no. 1, 1–13.
- [5] Hohlov, E. Y., *Operators and operations in the class of univalent functions*, Izv. Vyssh. Uchebn. Zaved. Mat., (1978), no. 10 (197), 83–89 (in Russian).
- [6] Jung, I. B., Kim, Y. C. and Srivastava, H. M., *The Hardy space of analytic functions associated with certain one-parameter families of integral operators*, J. Math. Anal. Appl. **176** (1993), no. 1, 138–147.
- [7] Kumar, V., Shukla, S. L., *Certain integrals for classes of  $p$ -valent meromorphic functions*, Bull. Austral. Math. Soc. **25** (1982), no. 1, 85–97.
- [8] Miller, S. S., *Differential inequalities and Carathéodory functions*, Bull. Amer. Math. Soc. **81** (1975), 79–81.
- [9] Noor, K. I., *On certain classes of meromorphic functions involving integral operators*, JIPAM. J. Inequal. Pure Appl. Math. **7** (2006), no. 4, Article 138, 8 pp. (electronic).
- [10] Noor, K. I., *On subclasses of close-to-convex functions of higher order*, Internat. J. Math. Math. Sci. **15** (1992), no. 2, 279–290.
- [11] Noor, K. I., *On new classes of integral operators*, J. Nat. Geom. **16** (1999), no. 1–2, 71–80.
- [12] Noor, K. I., *On close-to-convex and related functions*, Ph. D. Thesis, University of Wales, Swansea, U. K., 1972.
- [13] Noor, K. I., *On quasiconvex functions and related topics*, Internat. J. Math. Math. Sci. **10** (1987), no. 2, 241–258.
- [14] Noor, K. I., Noor, M. A., *On integral operators*, J. Math. Anal. Appl. **238** (1999), no. 2, 341–352.
- [15] Noor, K. I., Thomas, D. K., *Quasiconvex univalent functions*, Internat. J. Math. Math. Sci. **3** (1980), no. 2, 255–266.
- [16] Padmanabhan, K., Parvatham, R., *Properties of a class of functions with bounded boundary rotation*, Ann. Polon. Math. **31** (1975/76), no. 3, 311–323.
- [17] Pinchuk, B., *Functions of bounded boundary rotation*, Israel J. Math. **10** (1971), 6–16.
- [18] Ruscheweyh, S., *New criteria for univalent functions*, Proc. Amer. Math. Soc. **49** (1975), 109–115.
- [19] Selvaraj, C., Karthikeyan, K. R., *Some inclusion relationships for certain subclasses of meromorphic functions associated with a family of integral operators*, Acta Math. Univ. Comenian. (N. S.) **78** (2009), no. 2, 245–254.
- [20] Yuan, S.-M., Liu, Z.-M. and Srivastava, H. M., *Some inclusion relationships and integral-preserving properties of certain subclasses of meromorphic functions associated with a family of integral operators*, J. Math. Anal. Appl. **337** (2008), no.1, 505–515.

Ali Muhammad  
Department of Basic Sciences  
University of Engineering and Technology  
Peshawar  
Pakistan  
e-mail: [ali7887@gmail.com](mailto:ali7887@gmail.com)

Received March 25, 2011